

NEW HAMPSHIRE LAND COVER ASSESSMENT

Final Report

David Justice, Image Analyst
Anne Deely, Project Coordinator
Fay Rubin, GIS Manager

Complex Systems Research Center
University of New Hampshire
January, 2002



NEW HAMPSHIRE LAND COVER ASSESSMENT FINAL REPORT

DAVID JUSTICE, IMAGE ANALYST
ANNE DEELY, PROJECT COORDINATOR
FAY RUBIN, GIS MANAGER

COMPLEX SYSTEMS RESEARCH CENTER
UNIVERSITY OF NEW HAMPSHIRE
JANUARY, 2002

EXECUTIVE SUMMARY

Many organizations within the state and the region routinely need access to current and accurate information about New Hampshire's vegetative and physical features. In response to this need, the NH GRANIT staff at the Complex Systems Research Center, University of New Hampshire, conducted the New Hampshire Land Cover Assessment. The primary objective of the project was to generate a digital, statewide, 23-class land cover data set, developed from Landsat Thematic Mapper (TM) satellite imagery acquired over the period 1990-1999. Landsat 5 and 7 TM data were processed using a combination of traditional supervised and unsupervised classification methods. Data enhancements were achieved by utilizing supplemental sources archived in the GRANIT database, including Digital Orthophotoquads (DOQs), Digital Raster Graphics (DRGs), USGS Digital Line Graphs (DLGs) covering hydrography, NH Department of Transportation road centerlines, Digital Elevation Models (DEMs), SPOT panchromatic (10 meter resolution) images, public/conservation lands, US Fish & Wildlife Service National Wetlands Inventory (NWI) maps, and watershed boundaries.

A diverse coalition of agencies and organizations enabled the development of the land cover assessment through their financial support, cooperative field work, and data sharing. Field work conducted by project collaborators, including forestry professionals from the NH Department of Resources and Economic Development/Division of Forests and Lands, NH Fish and Game, and the U.S. Forest Service, was of great assistance, both in maintaining objectivity and in making the greatest use of limited resources. These organizations as well as the USDA Natural Resources Conservation Service, the Cooperative Institute for Coastal and Estuarine Environmental Technology (CICEET), and the NH Space Grant Consortium provided the financial support necessary to undertake the effort. The UNH Cooperative Extension provided assistance by sharing land management plans and supporting volunteer activities.

The product of the mapping effort is a digital land cover data layer covering the state of New Hampshire. The accuracy assessment indicates an overall accuracy of the aggregate, 7-class data

of 95.9%, and an accuracy of 82.2% at the full 23-class level. A supplemental product of the effort is a body of field data focused on the state's forested areas that provides a valuable data resource to a wide variety of users.

The land cover data set is available to resource managers, wildlife managers, foresters, educators, and the general public from the NH GRANIT website (www.granit.sr.unh.edu). It is also available on CD-ROM upon request.

Keywords: land cover, land use, land characterization, land resources, remote sensing, image processing.

INTRODUCTION

New Hampshire is the fastest growing state in the Northeast - its population has doubled since 1950, and will have tripled by 2020 (Sundquist and Stevens 1999). This population growth necessarily results in new development, including homes, schools, roads, and businesses, and the associated conversion of other land cover types (like forest and farmlands) to residential, commercial and industrial land uses. Accurate and timely data are needed in order to inventory and monitor these changing land use/land cover patterns.

It is well documented that satellite imagery, and specifically Landsat Thematic Mapper (TM) imagery, delivers a useful tool for regional land cover characterizations (Vogelmann, et al., 2001, Zhu, et al., 2000). It provides a data source for mapping large areas at a relatively low cost, and with a level of detail sufficient for many natural resource planning and management applications. Because it is acquired on a standard, 16-day cycle, it also provides a means to maintain and update land cover characterizations on a regular basis.

A number of previous efforts have utilized satellite imagery to produce land cover data for the state of New Hampshire. A project by researchers at the University of New Hampshire, and funded by the Environmental Protection Agency, was successful in deriving statewide land cover based on 1988-90 data sources (Rubin et al., 1993). A regional effort at the University of Vermont generated a broad land cover assessment for both New Hampshire and Vermont using 1992-93 imagery (Capen, et al., 2000). And at the federal level, the National Land Cover Data Set (Vogelmann, et al., 2001) used 1994-95 image sources to develop classified land cover data in generalized categories. Each of these products delivers useful data for trend analysis, but none is adequate in terms of data currency and/or classification detail to deliver the necessary data to state and local resource managers.

The New Hampshire Land Cover Assessment was undertaken to produce a current, statewide, digital land cover data set from 30-meter resolution Landsat 5 and 7 TM images. The imagery was processed to derive a 23-class land cover classification. The classification scheme was developed in collaboration with our project partners, and generally coincides with the USGS Anderson Level III scheme while providing a particular emphasis on the forested and agricultural classes (Table 1).

METHODS

Overview

Development of the statewide land cover data set progressed through a sequence of steps, from image acquisition and pre-processing through classification and accuracy assessment (Figure 1). The image classification phases included both traditional supervised and unsupervised techniques, which were supported by a body of training sites comprising newly-collected field data augmented by archived information. Data for the entire state were processed as a series of regional subsections corresponding to the geographic extents of the source images. Individual accuracy assessments were completed for each region, to assure satisfactory results at each processing stage. Finally, the regional classifications and accuracy assessments were compiled into a statewide data set.

Computer resources used throughout this project included the following hardware components: Silicon Graphics workstation platforms, a Dell Precision 420 workstation and a Dell Inspiron laptop computer. A number of software packages were utilized, including ERDAS Imagine, which was used for the bulk of the image processing tasks, and ESRI Arc/Info 7.x/8.x and ArcView 3.x, which were used for all phases of the geographic database management. For field navigation, we used Trimble PRO-XRS GPS receivers, each capable of receiving real time differential corrections (DGPS) via the U.S. Coast Guard beacon service and commercial satellite subscriptions. Trimble Pathfinder Office was used to complete global positioning tasks such as waypoint transfer and post-processing differential corrections (necessary when DGPS was not available).

Data Sources and Data Pre-processing

Twelve Landsat Thematic Mapper images, WRS path 12-13, row 29-30, spanning the years 1990-1999, were selected as the basis for the initial land cover classification (Table 2). The images were chosen to capture differing phenological conditions throughout the year. Each image was georeferenced to New Hampshire State Plane Coordinate feet (North American Datum of 1983, or NAD83) and terrain corrected. Georeferencing and terrain correction were performed either prior to the data purchase by the data vendors (USGS, EROS Data Center, or ImageLinks, Inc., Melbourne, FL) or post-purchase by UNH researchers. The images were then subset to comprise the geographic extent of three primary study areas, generally representing the southeast, southwest and northern areas of the state. Finally, the 6 reflective bands (1-5, and 7) from the most recent summer (leaf on) and spring (leaf off) image were “layer stacked,” or combined, into a single 12-band data set to be used in the initial, generalized classification for each region. The additional images were utilized in later processing stages.

Ancillary data comprised numerous holdings from the GRANIT archive, including panchromatic Digital Orthophotoquads (DOQs), Digital Raster Graphics (DRGs), USGS Digital Line Graphs (DLGs) covering hydrography, NH Department of Transportation road centerlines, Digital Elevation Models (DEMs), SPOT panchromatic (10 meter resolution) images,

public/conservation lands, US Fish & Wildlife Service National Wetlands Inventory (NWI) maps, and watershed boundaries (Figure 2).

Topographic Stratification

Previous efforts have suggested that preliminary topographic stratification can improve the accuracy of the results by offering the opportunity to isolate specific landscape characteristics (Rubin et al., 1993). Accordingly, topographic processing was initiated in the coastal region of the state where the 12-band, "layer-stacked" image was subset into three topographic categories: southeast facing slope (slope greater than 10% and azimuth between 45 - 225 degrees, clockwise), northwest facing slope (slope greater than 10% and azimuth between 225 - 45 degrees, clockwise) and flat (slope less than 10%). Large scale (1:24,000) USGS DEMs were used to isolate these three categories. Based on the preliminary coastal classifications, it was determined that using three slope categories in the remainder of the state would yield no appreciable increase in classification accuracy while requiring a significant increase in processing time and effort. Therefore, in the southwest and north regions, only two slope categories were subset: slope greater than 10%, and flat (slope less than 10%). Each aspect/slope stratum was then classified separately. Figure 1 presents a generalized diagram of the iterative classification process as applied to each aspect/slope category in each region. However, minor variations to this process may have occurred based on individual circumstances.

Training Sites Selection/Verification

In order to conduct supervised classifications, we utilized a body of training site data to represent the various targeted land cover classes. Some of these training sites were archived as a result of previous projects. However, a substantial amount of new data - over 1400 points - were collected to fill both regional and class-specific gaps, particularly in the forested and agricultural classes.

The image analyst identified candidate sites from the imagery based on its inherent characteristics. Typically, this involved observing color patterns and trying to account for the biological and spectral variability in the data. Additional forested sites, including beech/oak and hemlock, were selected based on DRED forest stand type maps. In each case, the image analyst identified points and used the ERDAS Imagine RegionGrow tool to select groups of spectrally similar pixels to form the training site. The resulting region in effect produced the representative signature used for the classifications. Generally, to be accepted as training sites, regions were required to contain 15 or more pixels, with a standard deviation across all image bands of 5 or less. Within the accepted pixel groupings, individual pixels were selected for subsequent verification.

In the case of the forested classes, 2-4 individual pixels were selected from each grouping for field sampling. The x-y coordinates of the pixel centroids were loaded into Trimble Pro-XRS GPS receivers (capable of real time differential correction, or DGPS, via Coast Guard beacon or satellite signal), which were then used to navigate to each training site point. In the rare case where DGPS was not available, the operator navigated to within 150 +/- feet of the point, collected position fixes, and corrected the position through standard post-processing. The

corrected point was then visually evaluated to determine if it was located within the training site polygon and was either accepted or discarded based on this position.

Once at the forested site, a 10 basal area factor (BAF) prism was used to tally all overstory trees within the variable radius plot. As the project developed, additional data were collected at each site for the southwest and northern regions (Figure 3). Ultimately, the prism tally was summarized and averaged by training site polygon, and the appropriate forest class label was calculated for each polygon according to the classification rules (Table 1). Field data collection for both the training site phase and the subsequent accuracy assessment was performed by GRANIT staff, assisted by professional foresters from the Division of Forests & Lands and the Fish & Game Department. Additionally, a corps of volunteers assisted the professional staff in the collection of field data in the southwest region (Figure 4).

Because the agricultural sites tended to be more homogeneous than the forested, a single point was sufficient for the field effort to identify the cover type. Again, GPS was used to navigate to the x-y coordinate, and the land cover was recorded. Additional information was not collected at these sites.

Table 3 reports the total number of field verified training sites available to the project for each land cover category, and Figure 5 shows the distribution of training site locations. The clustered nature of the data in the southern regions is a result of our attempt to restrict our activities to properties open to public access in order to avoid landowner permission issues. While relieving access issues, this approach also offered a secondary benefit as many of these properties had existing forest stand type information associated with them. This information was useful for targeting training sites to represent particular forested classes. In the north region, we received cooperation from several of the larger landowners through a direct liaison in the New Hampshire Fish and Game Department, who accompanied us to these forested training sites.

Data for a significant number of the non-forest and non-agricultural sites were amassed from pre-existing sources, such as DOQ's, DRG's, NWI, archived training sites from previous projects, and local knowledge, and did not require field work. For example, most of the training sites for cleared areas, such as disturbed and sand dunes, were developed from these sources. The wetland training sites were largely developed from the archived points and NWI data.

General Classification

A standardized procedure was utilized in each region of the state to produce a preliminary classification. First, a general classification was conducted by topographic stratum to group land cover types into five broad categories, including forested/coniferous, forested/deciduous, forested/mixed, agricultural/cleared, and wetlands/water. The processing began by applying archived training sites, primarily retrieved from past image classification projects, in order to create representative signatures. A supervised, maximum likelihood classification was then applied to the regional image, and the resulting classes were visually evaluated using DOQs, other ancillary data sets, and local knowledge. Acceptable classes were carried through to the final generalized data set, while unacceptable classes were retained for further processing (see Figure 1).

The unacceptable classes were subsequently used to mask various image band combinations and/or band transformations. (See Table 4a for a listing of bands/transformations used for each subset classification.) This step was followed by unsupervised classifications using the “isodata” cluster routine. At each iteration, the classes were evaluated as described above, and either archived for incorporation in the final product or retained for additional processing. As many as four supervised and unsupervised classification iterations using various image date/band derivatives were run on the resulting data subsets. Finally, each of the classifications was recoded to reflect the appropriate land cover value and mosaicked to generate the complete, region-wide generalized land cover data set.

Class-Specific Processing

With the generalized classification available, the analyst next addressed class-specific categories (e.g., wetland, forest, and agricultural types). Again, this was accomplished through a series of image subsets, masks and classification iterations to generate the final product. The additional images were introduced into the processing stream during this phase.

Each class-specific procedure was initiated by creating a “layer stack” of various image bands/band derivations, as reported in Table 4b. The bands were selected in part by applying the Imagine Signature Separability tool to the layer stack and using the Transformed Divergence measure as a basis for the selection. The tool reports metrics ranging from 0 - 2,000, with values equal to 2,000 reflecting signatures that are totally separable and values of 0 identifying signatures that are not separable. Once the appropriate bands were selected, the image composite was masked to retain pixels of interest (i.e., the forest specific classification retained forested classes from the generalized land cover). This was followed by an iterative process of classifications using a combination of techniques to derive the final data for that class.

The series of classifications typically began with a supervised classification, using both the archived training sites and the training sites collected for this project (Figure 5; Table 3). In the southeast, the three forested classes (conifer, deciduous, and mixed) from the generalized land cover were processed independently. However, since it was determined that there was no appreciable improvement in classification quality achieved by this separation, the three forested classes were processed together in the remaining regions.

As in the general classification, the class-specific scenario proceeded through a series of iterative classifications, with acceptable results saved for the final data layer and with unsatisfactory results used to mask subsequent data sets. For the forested classes, 2,794 training signatures were used to support the 14 iterations required to achieve an acceptable data layer. Note that some of these signatures were used redundantly; that is, the same training site was used to produce signatures for multiple images. For the cleared classes, 542 signatures were used while 12 classifications were run to develop an acceptable subset. And finally, 6 classifications and 126 signatures were implemented to complete the wetland data layer. (The wetland processing required many fewer training signatures due to our reliance on NWI data. We also utilized the Isodata routine more extensively for these classifications, which reduced the number of signatures necessary.)

Wetlands processing in the north region progressed in a slightly different manner than in the remainder of the state. After several classification attempts, it became clear that many forested wetlands, usually dominated by spruce/fir, were being confused with the upland spruce/fir class. While some classes from these processing iterations proved “usable” and thus were retained, many were not. Accordingly, we opted to use the NWI data as a mask to select the wetlands for class specific processing. All pixels coincident with the NWI data were classified into either the forested or open wetland classes. These were then added to “usable” wetland data set to create the entire wetland class for the north region.

Following each classification, either supervised or unsupervised, the image analyst evaluated each class to determine its “correctness.” Classes that were deemed accurate carried through to the final classification and poor classes were omitted. The analyst relied on ancillary data, such as NWI, DOQ, and DRG sources, as well as local knowledge, to determine the apparent reliability of each class. Ultimately, each of the “accepted” classified data layers was mosaicked into a single data set.

Finally, several post-processing refinements were applied to the provisional land cover data in the ESRI GRID environment. These enhancements included ancillary data overlay and data filtering. First, paved roads from the NH Department of Transportation road data (resident in the GRANIT data base, 2001) were “burned in” to the provisional land cover data set, effectively overwriting any coincident class. Secondly, DLG hydrography data were used to update double banked river, lake and pond edges. In many cases, this corrected problems in areas that were incorrectly mapped as conifer forest.

The forested classes were also modified based on existing topographic data. Elevation data from USGS digital elevation models were used to recode forest classes based on the U.S. Forest Service’s species elevation thresholds (Burns and Honkala 1990). Beech/oak above 2,500 feet and other hardwoods above 3,000 feet were recoded to paper birch/aspens; white/red pine above 1,500 feet and hemlock above 2,400 feet were recoded to spruce/fir; and any forested class above 4,200 feet was recoded to alpine (krumholz).

The fruit orchard class was also developed at this processing stage. Because past results indicated that spectral image classification of orchards was unlikely to yield acceptable results, orchard polygons were screen-digitized from DRG and DOQ images (where available), converted to the Arc/Info grid data structure and incorporated into the overall classification during post-processing.

Lastly, several filtering iterations were applied to the data set to remove speckling and produce minimum map units of one acre. In order to maintain the integrity of linear features, filtering was preceded by the RegionGroup command such that the majority filter would only operate on groups of pixels smaller than approximately 1 acre (5 pixels). This filter was next followed by a second RegionGroup and contiguous pixels fewer than 5 were finally Nibbled to remove those pixels that were not eliminated by the majority filter. This filtering processed the data set to its completed, final form.

Accuracy Assessment

A total of 975 sites were evaluated for the accuracy assessment. More than 600 of these sites were field visited, and others were evaluated using ancillary data such as NWI maps and DOQs. All sites classified as forest, and most classified as wetland, were field visited using Trimble ProXRS GPS units receiving real-time corrections. At forested sites, field crews recorded stand information and conducted up to five 10 BAF prism tallies to quantify stand composition. Figure 5 shows the geographic distribution of the assessment sites.

As with the classification itself, the accuracy assessment was conducted separately for each of the three geographic regions. In each region we attempted to sample 30 sites per land cover class, but this was not always achievable because of limited area covered by the class (such as with pitch pine), because of a post-data collection re-classification, or for other reasons. Conversely, some classes received higher than the targeted sampling levels, again because of the post-data collection re-classification. In order to limit distortion due to disparate sample sizes among classes, we randomly selected 20 sites from each class in each region to tabulate in the statewide error matrices. This yielded a total of 60 sites per class for the full error matrices (though some classes, particularly those like tundra that are regionally focused, still have fewer sample sites).

Error matrices were generated for the 23-class Level 3 classification, as well as the 17-class Level 2 and 7-class Level 1 aggregates. In each case, user's and producer's accuracies were calculated (Congalton and Green 1999). While these matrices provide valuable data, the traditional metrics alone did not appear to give potential users sufficient information on the magnitude and nature of errors. For example, a forested site containing a high percentage of beech and oak but tabulated to belong to the "mixed forest" class would be considered wrong in a traditional error matrix if classed as beech/oak. We felt it was important to identify broader ranges of "right" and "wrong" so that users could interpret the data most effectively. To address this, we created rules to assess the accuracy using fuzzy sets and the linguistic scale developed by Woodcock and Gopal (1992, 1994, 2000). The scale defines values ranging from 1 through 5 to indicate the extent to which an incorrect classification is "wrong", as follows:

- (1) *Absolutely wrong*: This answer is absolutely unacceptable. Very wrong.
- (2) *Understandable but wrong*: Not a good answer. There is something about the site that makes the answer understandable but there is clearly a better answer. This answer would pose a problem for users of the map. Not right.
- (3) *Reasonable or acceptable answer*: May not be the best possible answer but it is acceptable; this answer does not pose a problem to the user if it is seen on the map. Right.
- (4) *Good answer*: Would be happy to find this answer given on the map. Very right.
- (5) *Absolutely right*: No doubt about the match. Perfect.

Woodcock's and Gopal's rating of (2) was interpreted here to describe incorrect classifications that were understandable from an image processing perspective, but that would still be problematic for users. For example, the spectral characteristics of hay/pasture may be very similar to disturbed areas, and therefore it may be understandable that they are sometimes

confused in the classification, but these land uses are so different that this incorrect classification would be problematic to users of the data.

Table 5 contains the fuzzy set rules developed for this project. For forested sites, the rules for assigning a linguistic value to each sample site are quantitative, based on the proportions of basal area made up by different species. For non-forested sites, the rules are qualitative, based on how acceptable certain classifications may be at a given site. For example, for an open wetland, classification as open wetland would be absolutely right (5), classification as forested or tidal wetland would be a good answer (4), classification as open water would be understandable but wrong (2), and all other classifications would be absolutely wrong (1).

Each site was thus assigned a range of ratings based on the linguistic scale, and the rating that applied to the classification was identified. The accuracy of the map was then evaluated in terms of the how frequently the category assigned in the map was the best choice for the site (5), and how frequently the category assigned was acceptable (3-5) (Gopal and Woodcock 1994).

RESULTS

The primary result of this project is a 23-class, statewide digital land cover data layer (Figures 6 and 7). The data, archived in the GRANIT database, are accompanied by Federal Geographic Data Committee (FGDC)-compliant metadata. In addition, the project generated a body of field work that may be used for future analyses, including several thousand accurately positioned forested points with attributes describing species composition, size classes, and stand structure.

Table 6 presents acreage by land cover class and county for the entire state. As expected, the majority of the state (77.6% of total area, or 81.1% of land area) was mapped in the forested classes (not including forested wetlands) (Figure 8a). Grafton (84.7%) and Coos (84.3%) counties were the most heavily forested in the state, while Rockingham (58.6%) and Strafford (66.2%) were the least forested.

Deciduous forest was the most prominent forest class, covering approximately 32.9% (42.5% of total forest land) of the state. The majority of deciduous forest was either beech/oak or other hardwoods (Figure 8b). Mixed forest was also prominent, representing 26.3% of the state and 33.9% of the forestland. Finally, coniferous forests accounted for 18.2% of the state (23.6% of the total forest land), mostly composed of the white/red pine and spruce/fir classes.

Developed land was another class of significance, occupying 4.4% of the state. Table 6 shows that the most developed counties are Hillsborough (10.8%) and Rockingham (9.4%), and that the least developed is Coos county (1.1%). Slightly less of the state (4.1%) was mapped as agriculture. Most agricultural land (91.1% of total agricultural land) is in hay/pasture, while only 6.5% is in row crops and 2.4% is in fruit orchards. Grafton, Hillsborough, and Merrimack counties included the highest total acreage of agricultural land, while the least amount was mapped in Carroll and Strafford counties.

The classification achieved an overall accuracy of 82.2% at the full, 23-class level. Table 7 summarizes the results of the accuracy assessment in an error matrix, and shows user's and producer's accuracy percentages for each class. User's accuracy, or the percentage of occurrences in which a given map label correctly described the reference site, ranges from lows of 28.6% for paper birch/aspens, 53.3% for beech/oak, and 62.5% for mixed forest, to highs of 100% for water, tidal wetland, bedrock/vegetated, sand dune, and tundra. Producer's accuracy, or the percentage of occurrences in which reference sites were assigned a correct map label, ranges from lows of 28.6% for paper birch/aspens, 39.7% for mixed forest, 53.2% for other hardwoods, to highs of 100% for transportation, pitch pine, alpine, water, tidal wetland, bedrock/vegetated, sand dune, and tundra.

When collapsed to the 17-class level, the classification achieved an 88.7% overall accuracy, as shown in Table 8. Here too, there are broad ranges among the user's and producer's accuracies. Deciduous forest, whose components other hardwoods and paper birch/aspens fared relatively poorly at the third level classification, achieved a 94.8% user's accuracy and a 90.7% producer's accuracy. Coniferous forest, whose component hemlock fared relatively poorly at the third level classification, achieved an 81.9% user's accuracy and a 97.3% producer's accuracy.

At the aggregate, 7-class level, the classification achieved 95.9% overall accuracy (Table 9). User's accuracy ranged from 91.3% for developed to 100% for water and tundra, and producer's accuracy ranged from 89.9% for barren to 100% for water and tundra. Though the 23- and 17-class accuracy assessments show that it was difficult to distinguish between certain forested classes, the classification was consistently accurate in distinguishing forest from non-forest, achieving 99.0% producer's and 97.5% user's accuracy for forest.

Results of the fuzzy set assessment are summarized in Table 10. The "Max" and "% Max" columns correspond directly to the major diagonal and user's accuracy in the traditional error matrix; that is, they represent the number of sites that were correctly labeled on the map, and the percentage of sites with a given map label that were correctly validated by the reference data. The next columns expand the definition of correct from "max" (absolutely right, or those sites that were given a rating of 5) to include labels that were considered "right" (good answers, or those labels that were given ratings of 4 or 5) and, even more broadly, to those that were considered "reasonable or acceptable answers" (those labels that were given ratings of 3, 4, or 5).

The overall accuracy of the full classification increases to 89.1% when the "good answers" are included as "right," and to 92.0% when "reasonable or acceptable answers" are included as well. Particular improvement is seen in the forested categories - the accuracy of other beech/oak increases from 53.3% to 56.7% to 76.7%, and hemlock increases from 65% to 73.3% to 83.3%. Open wetland also improves dramatically, from 75% to 93.3%. Paper birch/aspens remains the least accurate class, due at least in part to its relatively small sample size.

DISCUSSION

The project successfully mapped the targeted land cover classes in New Hampshire. When used at an appropriate scale, users will have access to reliable data, with an overall accuracy

approaching 96% for the general classification and exceeding 82% for the 23-category classification. This product represents the most comprehensive and current land cover classification available for the region, and provides researchers with a reliable and well-documented data resource.

The standard accuracy assessment demonstrates a high degree of success in discriminating between deciduous and conifer classes. This was anticipated from the outset, due to the significant differences in the spectral properties evidenced by these two aggregate classes. The accuracies decline as more detail is introduced into the classification and species-level mapping is attempted, and particularly in the hardwood classes. There are several reasons for this decline in accuracy. In part, the difficulty may have resulted from our inability to acquire source imagery representing optimal phenology necessary to distinguish among the deciduous classes. Although Landsat satellites acquire data on a 16-day cycle, potentially yielding more than 20 scenes a year for a given area, in actuality, cloud cover and other sub-optimal conditions eliminate many of these. In fact, only eight usable October images were archived between the years of 1990 and 1998, and five of these were used in this project. In addition, though we expected that October images would provide the greatest deciduous phenological variation, the acquisition cycle is not always able to capture the short autumnal senescence.

The spatial resolution of the imagery also contributes to the difficulty of accurately delineating forest species classes. Landsat images are composed of pixels that are 30 meters square, and therefore can delineate features 900 m² (0.2 acres) or larger. (In fact, we have found that five-pixel blocks (one acre) are the minimum acceptable mapping unit from Landsat imagery.) However, the New Hampshire landscape seldom contains homogenous areas that large. Instead, the landscape, especially forested areas, contains a heterogeneous mix of land cover types. While single pixel blocks may be covered by forest (and accurately depicted as such through image classification), each block tends to be composed of a mosaic of different forest species classes covering much smaller areas, which cannot be captured using Landsat data.

There are also problems inherent in the targeted classification scheme (see Table 1). For example, the class definitions specify that a stand must be 75% deciduous to be labeled deciduous or 65% coniferous to be labeled coniferous. Therefore, a stand comprising 55% beech/oak and 45% hemlock would be incorrectly labeled if it were labeled as either of those classes. Instead, the “correct” label for this stand is mixed forest. To some degree, these problems are at least partially addressed using the fuzzy accuracy assessment. For example, researchers looking for hemlock stands may be disappointed to discover that the user’s accuracy is only 65%. If they are willing to accept forest stands with a minimum 50% hemlock basal area regardless of broad type (i.e., mixed forest), the class accuracy increases to 73.3%. Likewise, accepting a minimum hemlock basal area of 40% would yield a class accuracy of 83.3%. Not all classes make improvements of this magnitude with the fuzzy accuracy. Classes such as urban remain constant, while others, such as mixed forest, only marginally improve. However, we feel that the fuzzy set assessment provides a useful tool and that the relaxed requirements will be of interest to many data users.

The assessment of forested classes shows that the highest degree of success was achieved in mapping the white/red pine, spruce/fir and pitch pine classes. Each of these classes, and

especially pitch pine, has a very distinctive spectral signature. Mapping paper birch/aspen was particularly problematic. Both user's and producer's accuracies for this class are poor, demonstrating both that areas mapped as paper birch/aspen were actually other cover types, and that areas mapped as other cover types were actually paper birch/aspen. Most of the errors of commission reveal confusion with other deciduous classes; that is, sites mapped as paper birch were in fact other deciduous classes. This is probably a result of the spectral similarity between these classes. Most of the errors of omission reveal confusion with mixed forest; that is, sites mapped as mixed forest were in fact paper birch/aspen. This probably reveals that inadequate training sites were used for mixed forest and paper birch.

Mapping hemlock was also somewhat problematic. While there were relatively few errors in which sites with other map labels were in fact hemlock, there were many sites labeled as hemlock which were in fact other classes - most frequently, mixed forest. Because the accuracy improves considerably when we expand the classification rules to accommodate the fuzzy assessment, we see that much of the error is derived from the classification "cut-offs" between coniferous forest and mixed forest. Similarly, mixed forest was often labeled as paper birch/aspen, white pine, or other forest types. Again, this seems to stem from the difficulty inherent in creating discrete labels for continuously varying land cover classes. We also had difficulty distinguishing between beech/oak and other hardwoods. Aggregating these classes improves the accuracy dramatically, demonstrating that these classes were most often confused only with each other.

While most of the error in the forested wetlands classes is a result of confusion with open wetlands, in some cases, forest labels were misapplied to forested wetland sites. However, all such cases occurred in the coastal region. The classifications of the other regions of the state, which are based upon more recent imagery, show higher accuracies in the wetland classes, suggesting that age of source imagery may be one cause of the error in the wetlands categories. Beaver activity can rapidly convert forests to forested wetlands, and forested wetlands to open wetlands. In addition, open and forested wetlands are simply endpoints along a continuum, and it may be difficult for field crews to make an accurate visual interpretation of the wetland type.

The agricultural classes showed a high level of accuracy (95% producer's accuracy and 95.8% user's accuracy) at the generalized classification level (Table 9). As can be seen from the fuzzy assessment, the errors that did occur in these classes were most frequently a result of confusion between row crops and hay/pasture. Data sources quickly become outdated due to the dynamic nature of farming activities, and may have caused some of this confusion. Practices like crop rotation make it difficult to accurately classify agriculture sub-classes. However, some agricultural sub-classes – including orchards – are less variable. As we expected, the orchard mapping was quite accurate because of our use of DRGs in conjunction with DOQs (where available). Past experience has shown that this class would have been extremely difficult to map reliably based on spectral properties alone, as orchards commonly are classified as hay/pasture or other cleared.

Although we were generally satisfied with the overall accuracy of this product, some of the specific forested classes were troubling. It is our intention to revisit the data sources in an attempt to improve the accuracy of both the paper birch/aspen and mixed forested classes. It is

possible that a re-evaluation of the band combinations used in the classification of these categories may make it possible to improve their respective accuracies.

Also, we intend to continue our investigation into using radar imagery to predict relative forest densities. While the current classification maps broad forest types, little is known about the density of a given type (in this case expressed as basal area per acre). We will try to determine whether or not space borne radar imagery can assist in the mapping of forest densities in New Hampshire. It should be noted that the most appropriate radar data for this endeavor is at present not widely available, as the data we will use was acquired via the NASA Space Shuttle. However, with the launch of the NASDA ALOS satellite in 2003, L-band polarimetric radar data will become more commonly available. The results of this work will be discussed in an addendum to this report.

CONCLUSION

The New Hampshire Land Cover Assessment produced an accurate, statewide land cover data layer that provides resource managers, planners, researchers, and the general public with a creditable overview of New Hampshire's landscape. The land cover data set is available in ASCII grid format from the NH GRANIT website (www.granit.sr.unh.edu). It is also available on CD-ROM upon request. The data are accompanied by FGDC-compliant metadata, providing users with full documentation on the derivation of the data set, its technical characteristics, and recommendations on its appropriate use.

ACKNOWLEDGEMENTS

Partners:

NH Department of Fish and Game
NH Department of Resources and Economic Development
NH Space Grant
NOAA/UNH Cooperative Institute for Coastal and Estuarine Environmental Technology
UNH Cooperative Extension
USDA Forest Service
USDA Natural Resources Conservation Service

Landowners:

International Paper
Dartmouth College
Andorra Forest
Many other private landowners

Volunteers:

Liisa Adler
Dave Birchenough
Diane Fitzpatrick
Christine Lofgren
Andy Powell
Don Richard
Brian Rohde
Francie Von Mertens

Fieldwork:

David Justice, CSRC
Eric Swope, CSRC
Jennifer Merriam, CSRC
Ryan MacGregor, CSRC
Rusty Partridge, CSRC
Anne Deely, CSRC
Arin Mills, CSRC
Byard Mosher, CSRC
Lucie Plourde, CSRC
Jennifer Lessard, CSRC
Matt Gallucci, CSRC
Jennifer Bofinger, NH DRED
Garrett Dubois, NH DRED
Dave Falkenham, NH DRED
Kevin Finkenaur, NH DRED
Kyle Lombard, NH DRED
Robert MacGregor, NH DRED
Worthen Muzzey, NH DRED
Will Staats, NH Fish & Game
Kim Tuttle, NH Fish & Game
Tom Luther, US Forest Service

Project Support:

Bill Salas, CSRC
Susan Francher, DRED
Charlie Bridges, NH Fish and Game
John Lanier, NH Fish and Game
Nancy Lambert, UNH Cooperative Extension
Marshall Patmos, UNH Cooperative Extension
Tim Fleury, UNH Cooperative Extension
Steven Wood, UNH Cooperative Extension
Jon Nute, UNH Cooperative Extension
Norma Sorgman, US Forest Service

LITERATURE CITED

- Burns, R.M. and B.H. Honkala, technical coordinators. 1990. *Silvics of North America*. Agricultural Handbook 654, Volumes 1 and 2. U.S. Department of Agriculture, Forest Service, Washington, D.C., 1552 p.
- Capen, D.E., D.G. Williams, and E.W. Buford. 2000. Land cover map of Vermont & New Hampshire, 1993. <http://www.gap.uidaho.edu/>
- Congalton, R.G. and K. Green. 1999. *Assessing the accuracy of remotely sensed data: principles and practices*. Lewis Publishers, New York, 137 p.
- Gopal, S. and C. Woodcock. 1994. Theory and methods for accuracy assessment of thematic maps using fuzzy sets. *Photogrammetric Engineering and Remote Sensing* 60(2): 181-188.
- Rubin, F., D.G. Justice and J.E. Vogelmann. 1993. Final report: New Hampshire statewide digital wetlands inventory. Unpublished, 29 p.
- Sundquist, D. and M. Stevens. 1999. New Hampshire's changing landscape: population growth, land use conversion, and resource fragmentation in the Granite State. Society for the Protection of New Hampshire Forests and the New Hampshire Chapter of The Nature Conservancy, Concord, New Hampshire, 110 p.
- Vogelmann, J.E., S.M. Howard, L. Yang, C.R. Larson, B.K. Wylie, and N. van Driel. 2001. Completion of the 1990s National Land Cover Data Set for the conterminous United States from Landsat Thematic Mapper data and ancillary data sources. *Photogrammetric Engineering & Remote Sensing* 67(6): 650-662.
- Woodcock, C. and S. Gopal. 1992. Accuracy assessment of the Stanislaus Forest vegetation map using fuzzy sets. *Proceedings of the Fourth Biennial Remote Sensing Applications Conference*, Orlando, Florida, 17 p.
- Woodcock, C. and S. Gopal. 2000. Fuzzy set theory and thematic maps: accuracy assessment and area estimation. *International Journal of Geographical Information Science* 14(2): 153-172.
- Zhu, Z., L. Yang, S.V. Stehman, and R.L. Czaplewski. 2000. Accuracy assessment for the U.S. Geological Survey Regional Land-Cover Mapping Program: New York and New Jersey Region. *Photogrammetric Engineering & Remote Sensing* 66(12): 1425-1435.

Figure 1. Data Production Process.





Figure 2. Examples of data sources used for the classifications (Pawtuckaway Lake and environs, Nottingham, NH).

New Hampshire Land Cover Assessment




A. Landsat TM (1994)



B. SPOT Panchromatic (1993)



C. Digital Orthophotoquads (1992)



D. USGS Digital Raster Graphics

Figure 5. Locations of field sites, and approximate borders of classification regions. Field sites were visited outside the state to support the processing of data in adjoining study areas.




Figure 6. Level II statewide land cover map.




Figure 7. Level III land cover map of the Upper Contoocook watershed.




Figure 8. a) State area by land cover class. Total state area = 5,940,499 acres.
 b) Forest area by forest class. Total forested area = 4,607,287 acres.

New Hampshire Land Cover Assessment




Table 1. NH Land cover assessment classification scheme.

Level 1	Level 2	Level 3
1	Developed land	
	10 Residential/commercial/ industrial development	100 Residential/commercial/ industrial development
	14 Transportation	140 Transportation
2	Active agricultural land	
	21 Cropland and pasture	
		211 Row crops
		212 Hay/pasture
	22 Orchards, fruit, and ornamental horticulture	221 Orchards
4	Forest Areas dominated by trees, the majority of which are greater than 10' tall	
	41 Deciduous forest	Forest stands comprising less than 25% coniferous basal area per acre
		412 Beech/oak Deciduous stands comprising at least 30% beech and oak basal area per acre
		419 Other hardwoods All deciduous stands not meeting the beech/oak definition
	42 Coniferous forest	Forest stands comprising greater than 65% coniferous basal area per acre
		421 White/red pine Conifer stands in which white/red pine constitutes a plurality of the coniferous basal area
		422 Spruce/fir Conifer stands in which spruce/fir constitutes a plurality of the coniferous basal area
		423 Hemlock Conifer stands in which hemlock constitutes a plurality of the coniferous basal area
		424 Pitch pine Conifer stands in which pitch pine constitutes a plurality of the coniferous basal area

Table 1. NH Land cover assessment classification scheme.

Level 1		Level 2		Level 3	
		43	Mixed forest	430	Mixed forest Forest stands comprising more than 25% and less than 65% coniferous basal area per acre
		44	Alpine (krumholz)	440	Alpine (krumholz) Areas containing stunted vegetation, either hardwood or softwood, and occurring just below tree line in the White Mountains
5	Water	50	Open water	500	Open water Lakes, ponds, some rivers, or any other open water
6	Wetlands	Areas dominated by wetland characteristics defined by the U.S. Fish and Wildlife Service National Wetlands Inventory. Basically hydric soils, hydrophytic vegetation and the hydrologic conditions that result in water at or near the surface for extended periods of the growing season.			
		61	Forested wetlands	610	Forested wetlands Non-tidal wetlands characterized by woody vegetation 6m tall or higher
		62	Open wetlands	620	Open wetlands All other non-tidal wetlands, including those dominated by shrubs, emergents, mosses, or lichens
		63	Tidal wetlands	630	Tidal wetlands
7	Cleared/Other Open				
		71	Disturbed	710	Disturbed Gravel pits, quarries, or other areas where the earth and vegetation have been altered or exposed
		73	Sand dunes	730	Sand dunes Areas along the seacoast that are dominated by sand
		79	Other cleared	790	Other cleared Clear cut forest, old agricultural fields that are reverting to forest, etc.
8	Tundra	80	Tundra	800	Tundra Areas dominated by short vegetation and occurring above tree line on Mt. Washington

Table 2. Primary imagery data sources.

Image Type	Path	Row	Bands	Date	Source of Georeferencing/ Terrain Correction
Landsat 5 TM	12	30	1-7	8-Sep-90	CSRC
Landsat 5 TM	12	30	1-7	14-May-94	USGS
Landsat 5 TM	12	30	1-7	24-Oct-95	CSRC
Landsat 5 TM	12	30	1-7	22-Jul-96	USGS
Landsat 5 TM	13	29	1-7	13-May-91	USGS
Landsat 5 TM	13	29	1-5, 7	6-Oct-92	USGS
Landsat 5 TM	13	29	1-7	12-Oct-94	USGS
Landsat 7 ETM+	13	29	1-8	31-Aug-99	ImageLinks, Inc.
Landsat 5 TM	13	30	1-5, 7	6-Oct-92	USGS
Landsat 5 TM	13	30	1-7	28-Oct-94	USGS
Landsat 5 TM	13	30	1-7	14-Apr-98	USGS
Landsat 7 ETM+	13	30	1-8	31-Aug-99	ImageLinks, Inc.

Table 3. Field verified training sites available for the classifications.

Class	Southeast	Southwest	North	Archived	Total
Agriculture (aggregate)	0	0	0	66	66
Row crops	5	45	31	26	107
Hay/pasture	31	66	54	74	225
Beech/oak	214	125	87	142	568
Paper birch/aspen	32	21	25	14	92
Other hardwoods	58	75	47	245	425
Coniferous (aggregate)	5	0	0	133	138
White/red pine	38	66	27	23	154
Spruce/fir	0	43	25	7	75
Hemlock	49	29	10	5	93
Pitch pine	2	2	6	0	10
Mixed forest	67	77	19	101	264
Forested wetland	8	0	0	107	115
Open wetland	4	0	0	183	187
Disturbed land	0	0	1	13	14
Bedrock/vegetated	0	0	0	2	2
Other cleared	9	12	13	124	158
Total	522	560	345	1265	2692

Table 4. Bands and band transformations used for each subset classification.

a. Generalized Thematic Mapper classification

Subset	Path/Row	Image date	Raw TM Bands	PC ¹ Bands	TC ² Bands	NDVI ³	Band ratio
Southeast aspect	12/30	7/22/1996	1-5,7	1,2			
	12/30	10/24/1995		1,2			
	12/30	5/14/1994	1-5,7	1,2			
	12/30	9/8/1990		1,2			
	13/29	8/31/1999	1-5,7			1	
	13/29	10/12/1994	1-5,7		1,2,3		
	13/29	5/13/1991	1-5,7				
	13/30	8/31/1999					
	13/30	4/14/1998					
	13/30	10/28/1994					
	13/30	10/6/1992					
	13/30	8/31/1999	1-5,7				
	13/30	4/14/1998	1-5,7				
Northwest Aspect	12/30	7/22/1996	1-5,7				
	12/30	10/24/1995					
	12/30	5/14/1994	1-5,7				
	12/30	9/8/1990					
	13/29	8/31/1999	1-5,7			1	
	13/29	10/12/1994	1-5,7		1,2,3		
	13/29	5/13/1991	1-5,7				
	13/30	8/31/1999	1-5,7				
	13/30	4/14/1998	1-5,7				
Flat aspect	12/30	7/22/1996	1-5,7	1,2	1 - 6	1	
	12/30	10/24/1995	1				
	12/30	5/14/1994	1-5,7	1,2			5/4
	12/30	9/8/1990					
	13/29	8/31/1999	1-5,7	1,2,3			
	13/29	10/12/1994		1,2,3			
	13/29	5/13/1991	1-5,7	1,2,3			
	13/30	8/31/1999	1-5,7			1	
	13/30	4/14/1998	1-5,7		1,2,3		
	13/30	10/28/1994	3,4,5	1,2,3	1,2,3		
13/30	10/6/1992	3,4,5	1,2,3				

¹ Principal Components Analysis transformation

² Tasseled Cap Transformation

³ Normalized Difference Vegetation Index

Table 4. Bands and band transformations used for each subset classification.

b. Class-specific Thematic Mapper classification

Subset		Image date	Raw TM Bands	PC ¹ Bands	TC ² Bands	NDVI ³	Band ratio
Agriculture	12/30	7/22/1996	3-5, 7		1,2	1	
	12/30	10/24/1995			1,2	1	
	12/30	5/14/1994			1,2	1	
	12/30	9/8/1990			1,2	1	
	13/29	8/31/1999		1,2,3	1,2,3		
	13/29	10/12/1994		1,2			
	13/29	10/6/1992		1,2			
	13/29	5/13/1991		1,2			
	13/30	8/31/1999	1-5,7		1,2,3		
	13/30	4/14/1998	1-5,7				
	13/30	10/28/1994			1,2,3		
	13/30	10/6/1992	3,4,5		1,2,3		
Forest	12/30	7/22/1996		1-3		1	
	12/30	10/24/1995		1-3		1	
	12/30	5/14/1994		1-3		1	
	12/30	9/8/1990					
	13/29	8/31/1999		1,2,3			
	13/29	10/12/1994		1,2,3			
	13/29	10/6/1992		1,2,3			
	13/29	5/13/1991		1,2,3			
	13/30	8/31/1999		1,2,3			
	13/30	4/14/1998		1,2,3			
	13/30	10/28/1994		1,2,3			
	13/30	10/6/1992		1,2,3			
Wetland							
	12/30	7/22/1996	3-5, 7				
	12/30	10/24/1995					
	12/30	5/14/1994	3-5, 7				
	12/30	9/8/1990					
	13/29	8/31/1999	3-5,7				
	13/29	5/13/1991	3-5,7				
	13/30	8/31/1999			1-6	1	
	13/30	4/14/1998	1-5,7				
	13/30	10/28/1994	1-5,7			1	
13/30	10/6/1992	3,4,5			1		

¹ Principal Components Analysis transformation

² Tasseled Cap Transformation

³ Normalized Difference Vegetation Index

Table 5. Rules for the application of the fuzzy set linguistic scale.

Rating	
5	Absolutely right: No doubt about the match. Perfect.
4	Good answer: Would be happy to find this answer on the map.
3	Reasonable or acceptable: Not the best possible answer but acceptable; this answer does not pose a problem to the user if seen on the map.
2	Understandable but wrong: Not a good answer. There is something about the site that makes the answer understandable but there is clearly a better answer. This answer is problematic.
1	Absolutely wrong: This answer is absolutely unacceptable and completely wrong.

Beech/Oak (412):	
5	Deciduous stands comprising at least 30% beech and oak basal area per acre
4	Beech/oak BA greater than or equal to 30% regardless of broad type
3	Beech/oak BA greater than or equal to 15% and less than 30% regardless of broad type
2	Other forest
1	All other types

Other Hardwoods (419):	
5	All deciduous stands not meeting the beech/oak definition
4	All other deciduous stands
3	Mixed stands with softwood BA less than or equal to 35%
2	Other forest, row crops, hay/pasture, cleared other
1	All other types

White/Red Pine (421):	
5	Conifer stands in which white/red pine constitutes a plurality of the coniferous basal area
4	White/red pine BA greater than or equal to 50% regardless of broad type
3	White/red pine BA greater than or equal to 40% and less than 50% regardless of broad type
2	Other forest, forested wetlands
1	All other types

Spruce/Fir (422):	
5	Conifer stands in which spruce/fir constitutes a plurality of the coniferous basal area
4	Spruce/fir BA greater than or equal to 50% regardless of broad type
3	Spruce/fir BA greater than or equal to 40% and less than 50% regardless of broad type
2	Other forest, forested wetlands
1	All other types

Hemlock (423):	
5	Conifer stands in which hemlock constitutes a plurality of the coniferous basal area
4	Hemlock BA greater than or equal to 50% regardless of broad type
3	Hemlock basal area greater than or equal to 40% and less than 50% regardless of broad type
2	Other forest
1	All other types

Pitch Pine (424):	
5	Conifer stands in which pitch pine constitutes a plurality of the coniferous basal area
4	Pitch pine BA greater than or equal to 50% regardless of broad type
3	Pitch pine basal area greater than or equal to 40% and less than 50% regardless of broad type
2	Other forest, forested wetlands
1	All other types

Table 5. Rules for the application of the fuzzy set linguistic scale.

Mixed Forest (430):	
5	Forest stands comprising more than 25% and less than 65% coniferous basal area per acre
4	Coniferous BA less than 70% and greater than or equal to 65%; or coniferous BA greater than 20% and less than or equal to 25%
3	Coniferous BA less than 75% and greater than or equal to 70%; or coniferous BA greater than 15% and less than or equal to 20%
2	Other forest
1	All other types
Alpine (krumholz) (440):	
5	Alpine (krumholz)
4	Forest stands above 4200 feet in the White Mountains
3	None
2	None
1	All other types
Developed (100):	
5	Urban and built-up areas
4	Disturbed
3	None
2	None
1	All other types
Row crops (211):	
5	Row crops
4	Hay/pasture or orchards
3	None
2	Disturbed or other cleared (790) areas
1	All other types
Hay/Pasture (212):	
5	Hay/pasture
4	Row crops or orchards
3	Other cleared areas (790)
2	Urban and built up areas; disturbed areas
1	All other types
Orchards (221):	
5	Orchards
4	Row crops or hay/pasture
3	None
2	Other hardwood forest or mixed forest
1	All other types
Water (500):	
5	Open water
4	Non-forested wetlands or tidal wetlands
3	None
2	Forested wetlands
1	All other types

Table 5. Rules for the application of the fuzzy set linguistic scale.

Forested wetlands (610):	
5	Forested wetlands
4	Non-forested wetlands or tidal wetlands
3	None
2	Open water
1	All other types
Open wetlands (620):	
5	Non-forested wetlands
4	Forested wetlands or tidal wetlands
3	None
2	Open water
1	All other types
Tidal wetlands (630):	
5	Tidal wetlands
4	Forested or non-forested wetlands
3	None
2	Open water
1	All other types
Disturbed (710):	
5	Disturbed
4	None
3	Sand dunes or other cleared (790) areas
2	Row crops or hay/pasture
1	All other types
Sand dunes (730):	
5	Sand dunes
4	Disturbed
3	Cleared/other open
2	None
1	All other types
Other cleared (790):	
5	Other cleared (clearcuts, old fields, etc.)
4	Disturbed, row crops, or hay/pasture
3	Sand dunes or orchards
2	Deciduous forest
1	All other types
Tundra (800):	
5	Tundra
4	Bedrock/vegetated
3	Alpine (krumholz)
2	none
1	All other types

Table 6. Acreage summary by class and county.

Land Cover Class		County										New Hampshire
		Belknap	Carroll	Cheshire	Coos	Grafton	Hillsborough	Merrimack	Rockingham	Strafford	Sullivan	
Residential/ Commercial/ Industrial	Acres	4,704	5,348	6,956	4,002	9,434	28,137	13,397	14,210	4,344	4,556	95,089
	% of County	1.6%	0.8%	1.5%	0.3%	0.8%	4.9%	2.2%	3.1%	1.8%	1.3%	1.6%
Transportation	Acres	9,704	11,227	13,219	8,880	19,509	33,615	21,685	29,591	11,780	7,777	166,987
	% of County	3.2%	1.8%	2.8%	0.8%	1.7%	5.9%	3.5%	6.4%	4.8%	2.2%	2.8%
Total: Developed	Acres	14,408	16,576	20,175	12,882	28,943	61,752	35,082	43,801	16,124	12,333	262,077
	% of County	4.8%	2.6%	4.3%	1.1%	2.6%	10.8%	5.7%	9.4%	6.6%	3.5%	4.4%
Row Crops	Acres	396	305	2,397	1,934	3,344	1,065	3,266	506	503	2,167	15,882
	% of County	0.1%	0.0%	0.5%	0.2%	0.3%	0.2%	0.5%	0.1%	0.2%	0.6%	0.3%
Hay/ Pasture	Acres	13,075	10,468	24,212	19,058	40,643	31,418	34,190	19,636	10,535	20,676	223,909
	% of County	4.3%	1.6%	5.2%	1.6%	3.6%	5.5%	5.6%	4.2%	4.3%	5.9%	3.8%
Orchards	Acres	181	31	15	0	189	2,532	790	1,853	277	62	5,929
	% of County	0.1%	0.0%	0.0%	0.0%	0.0%	0.4%	0.1%	0.4%	0.1%	0.0%	0.1%
Total: Agriculture	Acres	13,652	10,804	26,624	20,992	44,175	35,014	38,246	21,995	11,314	22,904	245,719
	% of County	4.5%	1.7%	5.7%	1.8%	3.9%	6.1%	6.3%	4.7%	4.6%	6.5%	4.1%
Beech/ Oak	Acres	58,654	104,902	92,724	78,297	150,737	110,840	135,115	43,012	26,473	51,841	852,596
	% of County	19.5%	16.5%	19.9%	6.7%	13.5%	19.4%	22.1%	9.2%	10.8%	14.7%	14.4%
Birch/Aspen	Acres	1,242	38,570	8,137	82,021	99,229	5,603	4,934	242	0	6,474	246,453
	% of County	0.4%	6.1%	1.7%	7.0%	8.9%	1.0%	0.8%	0.1%	0.0%	1.8%	4.1%
Other Hardwoods	Acres	21,078	93,869	37,722	342,353	206,627	18,978	37,984	32,664	21,538	40,924	853,737
	% of County	7.0%	14.8%	8.1%	29.2%	18.5%	3.3%	6.2%	7.0%	8.8%	11.6%	14.4%

Land Cover Class		County										New Hampshire
		Belknap	Carroll	Cheshire	Coos	Grafton	Hillsborough	Merrimack	Rockingham	Strafford	Sullivan	
White/Red Pine	Acres	28,755	58,954	31,646	23,192	73,597	70,037	75,224	27,974	19,271	33,320	441,971
	% of County	9.6%	9.3%	6.8%	2.0%	6.6%	12.3%	12.3%	6.0%	7.9%	9.4%	7.4%
Spruce/Fir	Acres	3,467	35,914	15,833	187,646	122,496	9,967	16,982	136	41	32,346	424,828
	% of County	1.2%	5.6%	3.4%	16.0%	10.9%	1.7%	2.8%	0.0%	0.0%	9.2%	7.2%
Hemlock	Acres	9,165	24,835	33,373	12,457	45,542	26,442	30,858	4,944	1,955	20,563	210,133
	% of County	3.0%	3.9%	7.2%	1.1%	4.1%	4.6%	5.0%	1.1%	0.8%	5.8%	3.5%
Pitch Pine	Acres	14	3,714	0	0	0	3	31	16	9	0	3,788
	% of County	0.0%	0.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
Mixed Forest	Acres	79,360	146,623	154,309	255,453	246,430	159,019	165,211	163,596	92,730	100,761	1,563,493
	% of County	26.4%	23.1%	33.1%	21.8%	22.0%	27.8%	27.0%	35.2%	37.9%	28.5%	26.3%
Alpine (Krumholz)	Acres	0	0	0	6,619	3,670	0	0	0	0	0	10,289
	% of County	0.0%	0.0%	0.0%	0.6%	0.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%
Total: Forest	Acres	201,735	507,382	373,745	988,037	948,329	400,889	466,339	272,585	162,017	286,229	4,607,287
	% of County	67.1%	79.8%	80.1%	84.3%	84.7%	70.2%	76.3%	58.6%	66.2%	81.0%	77.6%
Open Water	Acres	48,461	45,081	17,367	26,641	29,351	15,529	19,734	32,882	15,985	11,680	262,711
	% of County	16.1%	7.1%	3.7%	2.3%	2.6%	2.7%	3.2%	7.1%	6.5%	3.3%	4.4%
Forested Wetland	Acres	1,498	9,031	3,023	20,835	4,963	4,711	3,994	12,385	3,706	1,133	65,278
	% of County	0.5%	1.4%	0.6%	1.8%	0.4%	0.8%	0.7%	2.7%	1.5%	0.3%	1.1%
Open Wetland	Acres	5,181	10,332	9,624	28,262	12,166	12,903	15,192	8,640	4,671	4,227	111,197
	% of County	1.7%	1.6%	2.1%	2.4%	1.1%	2.3%	2.5%	1.9%	1.9%	1.2%	1.9%
Tidal Wetland	Acres	0	0	0	0	0	0	0	5,120	115	0	5,235
	% of County	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.1%	0.0%	0.0%	0.1%

Land Cover Class		County										New Hampshire
		Belknap	Carroll	Cheshire	Coos	Grafton	Hillsborough	Merrimack	Rockingham	Strafford	Sullivan	
Total: Wetland	Acres	6,679	19,363	12,647	49,097	17,129	17,614	19,186	26,145	8,492	5,359	181,710
	% of County	2.2%	3.0%	2.7%	4.2%	1.5%	3.1%	3.1%	5.6%	3.5%	1.5%	3.1%
Disturbed	Acres	923	2,844	407	870	1,131	2,759	2,364	7,764	3,089	395	22,545
	% of County	0.3%	0.4%	0.1%	0.1%	0.1%	0.5%	0.4%	1.7%	1.3%	0.1%	0.4%
Bedrock/ Vegetated	Acres	0	921	283	434	962	13	33	0	0	568	3,215
	% of County	0.0%	0.1%	0.1%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.2%	0.1%
Sand Dunes	Acres	0	0	0	0	0	0	0	203	0	0	203
	% of County	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Other Cleared	Acres	14,930	32,828	15,271	70,021	49,484	37,586	30,164	59,793	27,844	13,888	351,808
	% of County	5.0%	5.2%	3.3%	6.0%	4.4%	6.6%	4.9%	12.9%	11.4%	3.9%	5.9%
Total: Cleared/ Other Open	Acres	15,853	36,593	15,961	71,326	51,576	40,358	32,560	67,760	30,933	14,851	377,772
	% of County	5.3%	5.8%	3.4%	6.1%	4.6%	7.1%	5.3%	14.6%	12.6%	4.2%	6.4%
Tundra	Acres	0	0	0	2,994	229	0	0	0	0	0	3,223
	% of County	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
Total	Acres	300,788	635,798	466,518	1,171,970	1,119,732	571,156	611,147	465,168	244,864	353,357	5,940,499

Table 7. Level 3 error matrix.

Land Cover Class		REFERENCE DATA													
		Resid./Comm./Indus.	Transportation	Row Crops	Hay/Pasture	Orchards	Beech/Oak	Paper Birch/Aspen	Other Hardwood	White/Red Pine	Spruce/Fir	Hemlock	Pitch Pine	Mixed Forest	
		110	140	211	212	221	412	414	419	421	422	423	424	430	
CLASSIFIED DATA	Resid./Comm./Indus.	110	53	0	0	0	0	0	0	0	0	0	0	0	
	Transportation	140	3	17	0	0	0	0	0	0	0	0	0	0	
	Row Crops	211	0	0	53	6	0	0	0	0	0	0	0	0	
	Hay/Pasture	212	0	0	1	55	1	0	0	0	0	0	0	0	
	Orchards	221	2	0	0	0	37	0	0	0	0	0	0	0	
	Beech/Oak	412	0	0	0	0	0	32	27	0	0	0	0	1	
	Paper Birch/ Aspen	414	0	0	0	0	2	4	5	0	0	0	0	3	
	Other Hardwood	419	0	0	0	0	13	2	42	0	0	0	0	1	
	White/Red Pine	421	0	0	0	0	0	0	0	49	0	1	0	9	
	Spruce/Fir	422	0	0	0	0	0	0	0	0	45	0	0	8	
	Hemlock	423	0	0	0	0	0	0	0	3	1	39	0	15	
	Pitch Pine	424	0	0	0	0	0	0	0	0	0	0	39	1	
	Mixed Forest	430	0	0	0	0	0	8	1	2	1	1	0	25	
	Alpine (Krumholz)	440	0	0	0	0	0	0	0	0	1	0	0	0	
	Water	500	0	0	0	0	0	0	0	0	0	0	0	0	
	Forested Wetland	610	0	0	0	0	0	0	2	0	0	0	0	0	
	Open Wetland	620	0	0	1	1	0	0	0	0	0	0	0	0	
	Tidal Wetland	630	0	0	0	0	0	0	0	0	0	0	0	0	
	Disturbed	710	3	0	1	1	0	0	0	0	0	0	0	0	
	Bedrock/ Veg.	720	0	0	0	0	0	0	0	0	0	0	0	0	
	Sand Dunes	730	0	0	0	0	0	0	0	0	0	0	0	0	
	Other Cleared	790	0	0	0	2	0	0	2	0	0	0	0	0	
	Tundra	800	0	0	0	0	0	0	0	0	0	0	0	0	
	Total		61	17	56	65	38	47	14	79	54	48	41	39	63

Table 7. Level 3 error matrix.

Land Cover Class		REFERENCE DATA										Total
		Alpine (Krumholz)	Water	Forested Wetland	Open Wetland	Tidal Wetland	Disturbed	Bedrock/ Veg.	Sand Dunes	Other Cleared	Tundra	
		440	500	610	620	630	710	720	730	790	800	
Resid./Comm./Indus.	110	0	0	0	0	0	5	0	0	2	0	60
Transportation	140	0	0	0	0	0	0	0	0	0	0	20
Row Crops	211	0	0	0	0	0	0	0	0	1	0	60
Hay/Pasture	212	0	0	0	0	0	0	0	0	3	0	60
Orchards	221	0	0	0	0	0	0	0	0	1	0	40
Beech/Oak	412	0	0	0	0	0	0	0	0	0	0	60
Paper Birch/ Aspen	414	0	0	0	0	0	0	0	0	0	0	14
Other Hardwood	419	0	0	1	1	0	0	0	0	0	0	60
White/Red Pine	421	0	0	1	0	0	0	0	0	0	0	60
Spruce/Fir	422	0	0	3	0	0	0	0	0	0	0	56
Hemlock	423	0	0	0	0	0	0	0	0	2	0	60
Pitch Pine	424	0	0	0	0	0	0	0	0	0	0	40
Mixed Forest	430	0	0	2	0	0	0	0	0	0	0	40
Alpine (Krumholz)	440	4	0	0	0	0	0	0	0	0	0	5
Water	500	0	20	0	0	0	0	0	0	0	0	20
Forested Wetland	610	0	0	52	5	0	0	0	0	1	0	60
Open Wetland	620	0	0	11	45	0	1	0	0	1	0	60
Tidal Wetland	630	0	0	0	0	20	0	0	0	0	0	20
Disturbed	710	0	0	0	0	0	54	0	0	1	0	60
Bedrock/ Veg.	720	0	0	0	0	0	0	20	0	0	0	20
Sand Dunes	730	0	0	0	0	0	0	0	20	0	0	20
Other Cleared	790	0	0	0	0	0	0	0	0	56	0	60
Tundra	800	0	0	0	0	0	0	0	0	0	20	20
Total		4	20	70	51	20	60	20	20	68	20	

Table 7. Level 3 error matrix.

Land Cover Class		Producer's Accuracy	User's Accuracy
Resid./Comm./Indus.	100	86.9%	88.3%
Transportation	140	100.0%	85.0%
Row Crops	211	94.6%	88.3%
Hay/Pasture	212	84.6%	91.7%
Orchards	221	97.4%	92.5%
Beech/Oak	412	68.1%	53.3%
Paper Birch/ Aspen	414	28.6%	28.6%
Other Hardwood	419	53.2%	70.0%
White/Red Pine	421	90.7%	81.7%
Spruce/Fir	422	93.8%	80.4%
Hemlock	423	95.1%	65.0%
Pitch Pine	424	100.0%	97.5%
Mixed Forest	430	39.7%	62.5%
Alpine (Krumholz)	440	100.0%	80.0%
Water	500	100.0%	100.0%
Forested Wetland	610	74.3%	86.7%
Open Wetland	620	88.2%	75.0%
Tidal Wetland	630	100.0%	100.0%
Disturbed	710	90.0%	90.0%
Bedrock/ Veg.	720	100.0%	100.0%
Sand Dunes	730	100.0%	100.0%
Other Cleared	790	82.4%	93.3%
Tundra	810	100.0%	100.0%
Overall Accuracy		82.2%	

Table 8. Level 2 error matrix.

Land Cover Class		REFERENCE DATA											
		Resid./Comm./Indus.	Transportation	Agriculture	Orchards	Deciduous Forest	Coniferous Forest	Mixed Forest	Alpine (Krumholz)	Water	Forested Wetland	Open Wetland	Tidal Wetland
		110	140	210	221	410	420	430	440	500	610	620	630
CLASSIFIED DATA	Resid./Comm./Indus.	110	53	0	0	0	0	0	0	0	0	0	0
	Transportation	140	3	17	0	0	0	0	0	0	0	0	0
	Agriculture	210	0	0	115	1	0	0	0	0	0	0	0
	Orchards	221	2	0	0	37	0	0	0	0	0	0	0
	Deciduous Forest	410	0	0	0	0	127	0	5	0	0	1	1
	Coniferous Forest	420	0	0	0	0	0	177	33	0	0	4	0
	Mixed Forest	430	0	0	0	0	9	4	25	0	0	2	0
	Alpine (Krumholz)	440	0	0	0	0	0	1	0	4	0	0	0
	Water	500	0	0	0	0	0	0	0	0	20	0	0
	Forested Wetland	610	0	0	0	0	2	0	0	0	0	52	5
	Open Wetland	620	0	0	2	0	0	0	0	0	0	11	45
	Tidal Wetland	630	0	0	0	0	0	0	0	0	0	0	20
	Disturbed	710	3	0	2	0	0	0	0	0	0	0	0
	Bedrock/ Veg.	720	0	0	0	0	0	0	0	0	0	0	0
	Sand Dunes	730	0	0	0	0	0	0	0	0	0	0	0
	Other Cleared	790	0	0	2	0	2	0	0	0	0	0	0
	Tundra	800	0	0	0	0	0	0	0	0	0	0	0
Total		61	17	121	38	140	182	63	4	20	70	51	20

Table 8. Level 2 error matrix.

CLASSIFIED DATA		REFERENCE DATA						Total			
		Disturbed	Bedrock/ Veg.	Sand Dunes	Other Cleared	Tundra					
		710	720	730	790	800					
Resid./Comm./Indus.	110	5	0	0	2	0	60	Resid./Comm./Indus.	110	86.9%	88.3%
Transportation	140	0	0	0	0	0	20	Transportation	140	100.0%	85.0%
Agriculture	210	0	0	0	4	0	120	Agriculture	210	95.0%	95.8%
Orchards	221	0	0	0	1	0	40	Orchards	221	97.4%	92.5%
Deciduous Forest	410	0	0	0	0	0	134	Deciduous Forest	410	90.7%	94.8%
Coniferous Forest	420	0	0	0	2	0	216	Coniferous Forest	420	97.3%	81.9%
Mixed Forest	430	0	0	0	0	0	40	Mixed Forest	430	39.7%	62.5%
Alpine (Krumholz)	440	0	0	0	0	0	5	Alpine (Krumholz)	440	100.0%	80.0%
Water	500	0	0	0	0	0	20	Water	500	100.0%	100.0%
Forested Wetland	610	0	0	0	1	0	60	Forested Wetland	610	74.3%	86.7%
Open Wetland	620	1	0	0	1	0	60	Open Wetland	620	88.2%	75.0%
Tidal Wetland	630	0	0	0	0	0	20	Tidal Wetland	630	100.0%	100.0%
Disturbed	710	54	0	0	1	0	60	Disturbed	710	90.0%	90.0%
Bedrock/ Veg.	720	0	20	0	0	0	20	Bedrock/ Veg.	720	100.0%	100.0%
Sand Dunes	730	0	0	20	0	0	20	Sand Dunes	730	100.0%	100.0%
Other Cleared	790	0	0	0	56	0	60	Other Cleared	790	82.4%	93.3%
Tundra	800	0	0	0	0	20	20	Tundra	810	100.0%	100.0%
Total		60	20	20	68	20		Overall Accuracy		88.4%	

Table 9. Level 1 error matrix.

CLASSIFIED DATA		REFERENCE DATA								Total
		Developed	Agriculture	Forest	Water	Wetland	Cleared/ Other Open	Tundra		
		100	200	400	500	600	700	800		
Developed	100	73	0	0	0	0	7	0	80	
Agriculture	200	2	153	0	0	0	5	0	160	
Forest	400	0	0	385	0	8	2	0	395	
Water	500	0	0	0	20	0	0	0	20	
Wetland	600	0	2	2	0	133	3	0	140	
Cleared/Other Open	700	3	4	2	0	0	151	0	160	
Tundra	800	0	0	0	0	0	0	20	20	
Total		78	159	389	20	141	168	20		

Land Cover Class		Producer's Accuracy	User's Accuracy
Developed	100	93.6%	91.3%
All Agriculture	200	96.2%	95.6%
Forest	400	99.0%	97.5%
Water	500	100.0%	100.0%
Wetland	600	94.3%	95.0%
Cleared/Other Open	700	89.9%	94.4%
Tundra	800	100.0%	100.0%
Overall Accuracy		95.9%	

Table 10. Results of the fuzzy logic assessment.

Map Label	"Absolutely right"		"Good Answers"		"Reasonable or Acceptable Answers"		Total
	Max (5)	% Max	Right (4-5)	% Right	Right (3-5)	% Right	
Residential/Commercial/Industrial	53	88.3%	58	96.7%	58	96.7%	60
Transportation	17	85.0%	20	100.0%	20	100.0%	20
Row Crops	53	88.3%	59	98.3%	59	98.3%	60
Hay/Pasture	55	91.7%	57	95.0%	60	100.0%	60
Orchards	37	92.5%	37	92.5%	37	92.5%	40
Beech/Oak	32	53.3%	34	56.7%	46	76.7%	60
Paper Birch/Aspen	4	28.6%	5	35.7%	5	35.7%	14
Other Hardwood	42	70.0%	57	95.0%	58	96.7%	60
White/Red Pine	49	81.7%	52	86.7%	53	88.3%	60
Spruce/Fir	45	80.4%	49	87.5%	52	92.9%	56
Hemlock	39	65.0%	44	73.3%	50	83.3%	60
Pitch Pine	39	97.5%	40	100.0%	40	100.0%	40
Mixed Forest	25	62.5%	27	67.5%	28	70.0%	40
Alpine (Krumholz)	4	80.0%	5	100.0%	5	100.0%	5
Water	20	100.0%	20	100.0%	20	100.0%	20
Forested Wetland	52	86.7%	57	95.0%	57	95.0%	60
Open Wetland	45	75.0%	56	93.3%	56	93.3%	60
Tidal Wetland	20	100.0%	20	100.0%	20	100.0%	20
Disturbed	54	90.0%	54	90.0%	55	91.7%	60
Bedrock/ Veg.	20	100.0%	20	100.0%	20	100.0%	20
Sand Dunes	20	100.0%	20	100.0%	20	100.0%	20
Other Cleared	56	93.3%	58	96.7%	58	96.7%	60
Tundra	20	100.0%	20	100.0%	20	100.0%	20
Total	801	82.2%	869	89.1%	897	92.0%	975